

HAI YANG¹, WENCHANG ZHOU¹, LIPING YANG², JIYONG DENG¹,
QIAN HU¹, HONGQING WEI¹, CHAKTONG AU¹, BING YI¹

Ag⁺/S₂O₈²⁻ SYSTEM FOR THE DEGRADATION OF AQUEOUS FLUTRIAFOL

The degradation of flutriafol in aqueous persulfate (S₂O₈²⁻) system in the presence of selected transition metal ions was investigated. In the presence of Ag⁺ or Fe²⁺, flutriafol degradation occurs, whereas in the presence of Mn²⁺, Co²⁺, Ni²⁺, Cu²⁺, Zn²⁺ and Bi³⁺ ions, the degradation is less than 15%. The Ag⁺/S₂O₈²⁻ aqueous system being the most effective was adopted for experimental and theoretical investigations. It was confirmed that pH of 3.0 is the most suitable, and that both [•]SO₄⁻ and [•]OH radicals are the main active species to afford flutriafol degradation, with the former contributes more than the latter. To elucidate degradation mechanism, molecular orbital calculations were performed and reaction intermediates identified by GC/MS and HPLC/MS/MS analyses. Three degradation pathways are proposed that involve the cleavage of C–N and C–C bonds as a result of [•]SO₄⁻ attack, as well as the formation of hydroxylated products due to [•]OH radicals.

1. INTRODUCTION

As a triazole fungicide (Fig. 1), flutriafol is used for systemic foliar or seed treatment to control plant diseases through the inhabitation of C-14 α -demethylase enzyme that participates in the biosynthesis of fungal sterols [1]. The compound has an important position in the global fungicide market because it can effectively control a vast number of diseases (e.g., cloud disease, leaf spot disease and rust disease) that affect a wide range of crops [2]. However, flutriafol is extremely persistent in soil, and being highly mobile, it is listed as a groundwater contaminant [3]. Despite not being used in the period of sampling, flutriafol at μg levels was detected in public water supply in England [4], Norway [5], Ireland [3] and other countries [6], indicating the persistence

¹College of Chemistry and Chemical Engineering, Hunan Institute of Engineering, Xiangtan 411104, P.R. China, corresponding author H. Yang, e-mail address: yanghai1001@163.com

²Shenzhen Academy of Environmental Sciences, Shenzhen 518001, P.R. China.

and accumulation of flutriafol in soil and ground water [7]. To understand the potential risk of secondary poisoning, it is meaningful to study the transformation of flutriafol in the environment. Flutriafol degrades through photolysis [8], oxidative [9], and biotransformation [10] processes. Nonetheless, the degradation rates are low under nutrient-limited conditions such as in subsurface soil or groundwater. For environment remediation, it is necessary to develop an effective method to determine the bioactivity, toxicity, metabolism, and degradation behavior of flutriafol in oxygen-depleted situations.




Fig. 1. Chemical structure of flutriafol with atoms labelled

Hydroxyl ($\cdot\text{OH}$) and sulfate ($\cdot\text{SO}_4^-$) radicals play important and decisive roles in the transformation of organic pollutants [11]. They both can initiate a series of chain reactions through which organic compounds are partially and/or fully decomposed [12]. Compared to $\cdot\text{OH}$ ($E^0 = 2.80$ V vs. NHE, $t_{1/2} = 10^{-3}$ μs), $\cdot\text{SO}_4^-$ ($E^0 = 2.43\text{--}2.80$ V vs. NHE, $t_{1/2} = 1\text{--}10$ μs) is similar in redox potential but more stable. There are $\cdot\text{SO}_4^-$ radicals in ground waters originating from atmospheric SO_2 , natural sulfur, and sulfur-based inorganic salts in industrial and household wastes [13].

Organic pollutants in water can be effectively degraded and transformed by $\cdot\text{SO}_4^-$. However, the ecotoxicology, metabolic transformation and degradation products of flutriafol in the environment cannot be monitored easily because of the low concentration and interference from inorganic/organic matters. Zhang et al. [13] selected the sulfate radical-advanced oxidation process (SR-AOP) to study the transformation of pharmaceuticals, which is a promising approach for soil and groundwater remediation [14]. Among various oxidants available, persulfate ($\text{S}_2\text{O}_8^{2-}$) is the newest and least studied, but has received increasing attention [15]. Once $\text{S}_2\text{O}_8^{2-}$ is activated by heat, UV light, bases, transition metals, soil minerals, and radiolysis, $\cdot\text{SO}_4^-$ radicals are formed [16]. In soil and groundwater, $\cdot\text{SO}_4^-$ reacts with a wide variety of contaminants, generating reactive species such as $\cdot\text{OH}$, HSO_5^- and H_2O_2 [17].

The $\text{S}_2\text{O}_8^{2-}$ -based SR-AOP is superior in removing organic pollutants. Romero et al. [18] reported the degradation of diuron in a $\text{Fe(II)}/\text{S}_2\text{O}_8^{2-}$ system, and achieved complete oxidation of diuron in few minutes at 0.72 mmol/dm^3 Fe(II) concentration. The same

Fe(II)/S₂O₈²⁻ system was chosen by Zhou et al. [19] to remove diuron and As(III), and [•]OH and [•]SO₄⁻ are key participants in the oxidation process in which As(III) is oxidized to As(V) via an electron-transfer step. Adopting the heat/S₂O₈²⁻ system, Tan et al. [20] studied the degradation of diuron, and reported that [•]SO₄⁻ was the principal agent responsible for its degradation. Furthermore, Mendez-Diaz et al. [21] examined the degradation efficiency of surfactant SDBS, and found that the UV/S₂O₈²⁻ process was suitable. Despite these successful cases, the transformation mechanism of organic pollutants in S₂O₈²⁻-based SR-AOP are not fully recognized. Moreover, it is a challenge to explore feasible methods for efficient generation of [•]SO₄⁻ from S₂O₈²⁻.

In the present study, the degradation of flutriafol in persulfate-based SR-AOPs was investigated theoretically and experimentally. First the S₂O₈²⁻ activation efficiencies of selected transition metals were compared. Since silver was found to be the most efficient, the Ag⁺/S₂O₈²⁻ system was adopted to investigate the influence of chelating agents, pH values and substrate concentrations on flutriafol degradation. The related transformation mechanism was elucidated through molecular orbital calculations and identification of degradation intermediates.

2. MATERIALS AND METHODS

Materials. Flutriafol (≥99% purity) was purchased from the Hunan Research Institute of Chemical Industry. Sodium persulfate (Na₂S₂O₈, 99.1%) and ferrous sulfate heptahydrate (FeSO₄·7H₂O, 98%) were purchased from Sigma-Aldrich. Transition metal compounds (MnCl₂, Ni(NO₃)₂·6H₂O, ZnCl₂, Co(NO₃)₂·6H₂O, AgNO₃, Cu(NO₃)₂, Bi(NO₃)₃·5H₂O and FeCl₂) and the metal ion chelators: SCD (98%), Na₂S₂O₃·5H₂O (98%), DM (99%), and EDTA-Na₂ (98%), were supplied from Shanghai Sinopharm Chemical Reagent Co., Ltd., China. HPLC grade water was generated using a Millipore Milli-Q system in which the total organic carbon concentration was kept below 13 μg/dm³ by constant illumination with a xenon arc lamp at 172 nm. Acetonitrile, menthol and ethanol (HPLC grade) were obtained from Sigma. The reagents were AR grade unless mentioned otherwise.

Experimental procedures. Homogeneous reactions were run at ambient temperature (25±2 °C) in a 250 cm³ glass reactor wrapped with aluminum foil. A specific aliquot of 25–200 μmol/dm³ flutriafol was transferred into the reactor, followed by the addition of a designated amount of Na₂S₂O₈ (0.1–10 mmol/dm³) and transition metal salts (1.0 mmol/dm³ MnCl₂, Ni(NO₃)₂·6H₂O, ZnCl₂, Co(NO₃)₂·6H₂O, AgNO₃, Cu(NO₃)₂, Bi(NO₃)₃·5H₂O and FeCl₂). The initial pH (1–11) of the solution was adjusted by using 0.1 mol/dm³ HCl and 0.1 mol/dm³ NaOH. The solution was constantly stirred with a magnetic stirrer. 1 cm³ samples of the solution were taken at fixed time intervals, and quenched immediately by 1 cm³ menthol, a quenching agent for [•]OH and [•]SO₄⁻ radicals,

and then analyzed (after going through a 0.22 μm Millipore filter) by the HPLC and HPLC/MS/MS techniques. In the experiments of chelating agents, 1 mmol/dm^3 SCD, EDTA- Na_2 , $\text{Na}_2\text{S}_2\text{O}_3$ and DM were added to simulate interference in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system ($\text{S}_2\text{O}_8^{2-}$ and Ag^+ concentration – 2.0 and 1.0 mmol/dm^3 , respectively). Similarly, 0.1 mol/dm^3 ethanol and NaHCO_3 were added to $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system in the experiments of reactive oxygen species (ROS) scavengers, respectively. For GC/MS analysis, $50\times 3\text{ cm}^3$ CH_2Cl_2 was added to the sample solution for extraction. The organic phase was collected, dried over Na_2SO_4 to desolventize at ambient temperature. The degradation efficiencies were calculated from the formula $((C_0 - C)/C_0\times 100\%)$. Each batch of experiments was performed in triplicate, and the deviation was less than 5% in most of the cases.

Analytic methods. Flutriafol analysis was performed at 23 $^\circ\text{C}$ over a Cometro 6000 series HPLC using Kromasil C18 column ($250\times 4.6\text{ mm}$). The mobile phase was composed of 40% H_2O and 60% CH_3CN that was filtered with a Water Associates (Milford; MA, USA) 0.45 μm filter. The detection wavelength was 220 nm. The flow rate of the mobile phase was set at 1 cm^3/min .

Chromatographic analysis was carried out over a Shimadzu GCMS-QP2010 gas chromatograph/mass spectrometer equipped with a fused-silica capillary column coated with 5% diphenylmethylsiloxane, viz. VF-5 ($30\text{ m}\times 0.25\text{ mm}$, 0.25 μm film thickness) from Varian. High-purity helium (99.9999%) at a constant flow rate of 1.5 cm^3/min was used as carrier gas. Employing an AOC-20i auto injector (275 $^\circ\text{C}$), injections of 10^{-3} cm^3 were made in the splitless mode with 1.0 min purge-off time. The temperature program for analysis was: initial temperature 60 $^\circ\text{C}$ (held for 2 min), temperature raised (10 $^\circ\text{C}/\text{min}$) to 200 $^\circ\text{C}$ (held for 1 min), and then to 275 $^\circ\text{C}$ and held at this temperature for 10 min. The GC was directly interfaced with the Shimadzu QP 2010 quadrupole mass spectrometer (250 $^\circ\text{C}$ interface temperature and 70 eV impact ionization). The transfer line was set at 275 $^\circ\text{C}$, and the source at 200 $^\circ\text{C}$. Positive fragment ions were analyzed over the 43–500 m/z mass range in SCAN mode. Unequivocal identification of target compounds was achieved through comparison with mass-spectral and special pesticide libraries (e.g., NIST, Wiley, PEST).

Degradation intermediates were analyzed using a HPLC/MS/MS equipment. HPLC separation was performed at 0.5 cm^3/min mobile phase (50% of H_2O and 50% of CH_3CN) within 30 min over a Shimadzu HPLC system equipped with a Kromasil C18 column ($150\times 4.6\text{ mm}$), and SIL-HT autosampler. With LC-10 AT vacuum pump and API 3000 mass analyzer, and through electrospray interface, MS and MS-MS analysis in positive ionization mode for full scan acquisition between m/z 50–450 were performed. The collision energy was varied according to measurement requirement, and the other parameters were set as follows: ESI 5.5 keV, source block and desolvation temperature 130 $^\circ\text{C}$ and 400 $^\circ\text{C}$, respectively, desolvation and nebulizer gas (N_2) flow rate 6 dm^3/min , and argon (as collision gas) pressure 250 kPa.

Molecular orbital calculations were performed using Gaussian 03 program (Gaussian, Inc.) at the single determinant (HF/3-21) level. The optimal conformation having a minimum energy was obtained at the B3LYP/6-31G* level. The $(FED_{HOMO}^2 + FED_{LUMO}^2)$ and $2FED_{HOMO}^2$ values were acquired to predict the initial state of SO_4^- and OH interaction with the flutriafol molecule, respectively [22]. Also, the PCs were calculated based on NBO theory.

3. RESULTS AND DISCUSSION

3.1. $S_2O_8^{2-}$ ACTIVATION BY TRANSITION METALS

The effect of Mn^{2+} , Ni^{2+} , Zn^{2+} , Co^{2+} , Ag^+ , Cu^{2+} , Bi^{3+} and Fe^{2+} on persulfate activation was studied (Fig. 2). For the investigation, 1.0 mmol/dm^3 of metal cation was added, and NO_3^- or Cl^- was the counter anion.




Fig. 2. Degradation efficiency of $100 \text{ } \mu\text{mol/dm}^3$ flutriafol in the 1.0 mmol/dm^3 $S_2O_8^{2-}$ system activated by various transition metal ions

It was found that flutriafol ($100 \text{ } \mu\text{mol/dm}^3$) degradation is 100% in 120 min with Ag^+ whereas it is less than 15% in 360 min with Mn^{2+} , Ni^{2+} , Zn^{2+} , Co^{2+} , Cu^{2+} and Bi^{3+} . With Fe^{2+} , flutriafol degradation is 82% in 360 min. Among the eight cases, flutriafol degradation is the worst with Bi^{3+} but best with Ag^+ . The poor performance of Bi^{3+} is ascribed to its saturated electron configuration which is $5s^25p^65d^{10}6s^2$. With such electron configuration, Bi^{3+} is chemically inert. The electron configurations of Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} and Zn^{2+} are $3s^23p^63d^5$, $3s^23p^63d^6$, $3s^23p^63d^7$, $3s^23p^63d^8$, $3s^23p^63d^9$, and

$3s^23p^63d^{10}$, respectively. Despite the metal ions can accept electrons before attaining saturated electron configuration, they are poor in persulfate activation with Fe^{2+} being the only exception. It is because the $Metal^{3+}/Metal^{2+}$ potentials of Mn^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} and Zn^{2+} are high while that of Fe^{2+} (Fe^{3+}/Fe^{2+} , $E^0 = 0.77$ V vs. NHE) is relatively low [22]. Interestingly, flutriafol degradation is the most efficient in the $Ag^+/S_2O_8^{2-}$ system. Ag^+ has an extra-nuclear electron configuration of $4s^24p^64d^{10}$ that allows the loss of an electron in the presence of a strong oxidant such as $S_2O_8^{2-}$. The standard Ag^{2+}/Ag^+ potential ($E^0 = 1.98$ V vs. NHE) is lower than the $S_2O_8^{2-}/SO_4^{2-}$ ($E^0 = 2.01$ V vs. NHE) and SO_4^{2-}/SO_4^- ($E^0 = 2.43-2.80$ V vs. NHE) potential, indicating that the transformation of Ag^+ to Ag^{2+} is feasible in the $Ag^+/S_2O_8^{2-}$ system.

3.2. SINGLE VARIABLE AT A TIME EXPERIMENTS IN $Ag^+/S_2O_8^{2-}$ SYSTEM

The flutriafol degradation ability of Ag^+ or $S_2O_8^{2-}$ is low. As shown in Fig. 3, there is only insignificant degradation of flutriafol over each of them. It is only in the presence of Ag^+ that a stronger oxidative ability of $S_2O_8^{2-}$ is released. We investigated the effect of Ag^+ concentration on the degradation of $100 \mu M/dm^3$ flutriafol in the $Ag^+/S_2O_8^{2-}$ system by keeping $S_2O_8^{2-}$ constant at $2.0 \text{ mmol}/dm^3$. As shown in Fig. 4, an increase of Ag^+ concentration from 0.1 to $10 \text{ mmol}/dm^3$ results in gradual enhancement of degradation efficiency. And there is complete degradation of flutriafol in 10 min when Ag^+ concentration is $10 \text{ mmol}/dm^3$.




Fig. 3. Degradation efficiency of $100 \mu\text{mol}/dm^3$ flutriafol in $1.0 \text{ mmol}/dm^3$ Ag^+ , $S_2O_8^{2-}$ and $Ag^+/S_2O_8^{2-}$ systems




Fig. 4. Effect of Ag^+ concentration on the degradation efficiency of $100 \mu\text{mol/dm}^3$ flutriafol in the $Ag^+/S_2O_8^{2-}$ system; the $S_2O_8^{2-}$ concentration – 2.0 mmol/dm^3




Fig. 5. Effect of $S_2O_8^{2-}$ concentration on the degradation efficiency of $100 \mu\text{mol/dm}^3$ flutriafol in the $Ag^+/S_2O_8^{2-}$ system; the Ag^+ concentration – 1.0 mmol/dm^3

On the other hand, when Ag^+ concentration is kept constant at 1.0 mmol/dm^3 , there is increase of degradation efficiency when $S_2O_8^{2-}$ concentration is increased from 0.1 to 5.0 mmol/dm^3 (Fig. 5). And there was complete degradation of $100 \mu\text{mol/dm}^3$ flutriafol in around 30 min when $S_2O_8^{2-}$ concentration was 5.0 mmol/dm^3 . However, a further in-

crease of $S_2O_8^{2-}$ concentration to 10 mmol/dm^3 does not result in any increase of degradation efficiency. This might be due to the fact that too high $S_2O_8^{2-}$ concentration would result in other ROS being out-competed, leading to the stagnant of degradation efficiency [23].




Fig. 6. Degradation efficiency of $100 \text{ }\mu\text{mol/dm}^3$ flutriafol in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system with various chelating agents (1 mmol/dm^3), $\text{S}_2\text{O}_8^{2-}$ and Ag^+ concentrations 2.0 and 1.0 mmol/dm^3 , respectively




Fig. 7. Effect of flutriafol concentration ($25\text{--}200 \text{ }\mu\text{mol/dm}^3$) on the degradation efficiency of flutriafol in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system, $\text{S}_2\text{O}_8^{2-}$ and Ag^+ concentrations 2.0 and 1.0 mmol/dm^3 , respectively

The degradation of flutriafol was not affected by the addition of SCD, as seen in Fig. 6. However, with the addition of EDTA-Na₂ and DM, there is near complete suppression of flutriafol degradation, plausibly a result of their interaction with Ag⁺, i.e., capturing Ag⁺ through complex formation. With the capture of Ag⁺, the activation of S₂O₈²⁻ was denied. In the presence of Na₂S₂O₃, there was certain suppression of flutriafol degradation, which is a result of Ag⁺ interaction with S₂O₃²⁻, giving Ag₂S₂O₃ and Ag₂S (Eqs. (1) and (2)). The results confirm that once there is the capture of Ag⁺ ions, the ability of the Ag⁺/S₂O₈²⁻ system for flutriafol degradation weakens.



With the concentration of S₂O₈²⁻ and Ag⁺ fixed at 2 and 1 mmol/dm³, respectively, we studied how the change of flutriafol concentration would influence the degradation efficiency (Fig. 7). From a flutriafol concentration of 25–200 μmol/dm³, time for near complete degradation stretches from 10 to 180 min, indicating the amount of ROS in the Ag⁺/S₂O₈²⁻ system is limited.




Fig. 8. Effect of pH on degradation efficiency of 100 μmol/dm³ flutriafol in the Ag⁺/S₂O₈²⁻ system, S₂O₈²⁻ and Ag⁺ concentrations – 2.0 and 1.0 mmol/dm³, respectively

pH of the solution is an important parameter affecting oxidative degradation of organic contaminants. We found that a change of initial pH value (from 1 to 11) affects

the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system immensely (Fig. 8). When the solution is strongly acidic (pH 1) or neutral (pH 7), there is no degradation of flutriafol. It is only when the solution is weakly acidic (pH 3) or basic (pH 9 and 11) the degradation of flutriafol is observed. The lowest degradation efficiency at pH 1 is due to the removal of Ag^+ as a result of AgCl precipitation.



With the removal of Ag^+ , there is no activation of $\text{S}_2\text{O}_8^{2-}$. Nonetheless, we cannot give a proper explanation for the poor degradation efficiency at pH 7. Further experiments have to be performed to clarify such a phenomenon.

3.3. EFFECTS OF ROS AND PREDICTION OF ACTIVE SITES

To identify the dominating radical species in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system, quenching experiments were performed. Ethanol and NaHCO_3 were selected as quenching agents since the former is known to quench $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$ while the latter $\cdot\text{SO}_4^-$ [24, 25], and the results are shown in Fig. 9.




Fig. 9. Degradation curves of $100 \mu\text{mol}/\text{dm}^3$ flutriafol and flutriafol with scavengers ($0.1 \text{ mol}/\text{dm}^3$) in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system, $\text{S}_2\text{O}_8^{2-}$ and Ag^+ concentrations – to 2.0 and $1.0 \text{ mmol}/\text{dm}^3$, respectively

With the addition of $0.1 \text{ mol}/\text{dm}^3$ NaHCO_3 to the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system the degradation efficiency decreases to 57%, suggesting that 43% degradation is due to $\cdot\text{SO}_4^-$. When $0.1 \text{ mol}/\text{dm}^3$ ethanol is used, the degradation efficiency decreases significantly to 28%, suggesting that 72% degradation is due to $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$. A simple calculation shows

that the contribution of $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$ is 43% and 29%, respectively, and the rest 28% is due to ROS such as peroxysulfate.

Table 1

Point charges (PCs) and frontier electron densities (FEDs) on atoms of flutriafol calculated by Gaussian 03 program at the B3LYP/6-31G*

Atom label ^a	2FED _{HOMO} ²	FED _{HOMO} ² + FED _{LUMO} ²	PCs
C ₁	0.0896	0.0453	-0.0146
C ₂	0.2006	0.3231	0.4056
C ₃	0.0595	0.2017	-0.0150
C ₄	0.0790	0.0423	0.0408
C ₅	0.2815	0.3723	-0.0743
C ₆	0.0509	0.2527	0.1068
C ₇	0.0120	0.0136	0.2075
C ₈	0.0345	0.0706	0.4266
C ₉	0.1130	0.0928	-0.1281
C ₁₀	0.0015	0.0512	0.0147
C ₁₁	0.0782	0.0491	0.0031
C ₁₂	0.0643	0.0499	0.0277
C ₁₃	0.0047	0.0422	-0.0083
C ₁₄	0.0743	0.0448	0.4784
N ₁₅	0.0148	0.1148	-0.6673
N ₁₆	0.0002	0.0631	-0.3743
C ₁₇	0.0009	0.0312	0.5726
N ₁₈	0.0019	0.0113	-0.6287
C ₁₉	0.0007	0.0864	0.6985
F ₂₀	0.0813	0.0616	-0.4105
O ₂₁	0.0093	0.0182	-0.2683
F ₂₂	0.0242	0.0126	-0.3931

^aFor the label of atoms refer to Fig. 1.

According to the frontier orbital theory, we looked into the nature of the active sites in the interaction of the flutriafol molecule with ROS. It is known that the locations with higher FED_{HOMO}² + FED_{LUMO}² values show higher tendency to interact with radicals, while those with higher 2FED_{HOMO}² values are likely to lose electrons in oxidation reaction [26]. As shown in Table 1, the C₂, C₃, C₅ and C₆ of flutriafol show high FED_{HOMO}² + FED_{LUMO}² values of 0.4457, 0.3438, 0.4632 and 0.2527, respectively. The two benzene rings are different in reaction activity. The F atom at the *para*-position activates the benzene ring while that at the *ortho* position deactivates the benzene ring, resulting in the former being more reactive towards radical interaction. On the other hand, C₂, C₅ and C₉ show higher 2FED_{HOMO}² values, indicating that these atoms of the benzene rings

are likely to lose electrons and get involved in oxidation reaction. As for C₂ and C₅, reactivity is low due to steric effect. It is hence reckoned that C₉ was the predicted position for oxidation reaction at the initial stage. Furthermore, the PC values show that C₂, C₈, C₁₄, C₁₇ and C₁₉ are positive while the heteroatoms N, O and F atoms are negative. Accordingly, the carbon atoms in the triazole ring and those connected with the F atoms have tendency to react with radicals, while the heteroatoms of the triazole ring are likely to react with H⁺ and Ag⁺.

3.4. INVESTIGATION OF DEGRADATION INTERMEDIATES AND TRANSFORMATION MECHANISM

By means of GC/MS and HPLC/MS/MS analysis we investigated the degradation intermediates and the transformation mechanism, and the TIC chromatograms of GC/MS and HPLC/MS/MS are presented in Figs. 10 and 11, respectively.




Fig. 10. TIC chromatogram from GC/MS

From Fig. 10, one can see five degradation intermediates, viz. P₁-140, P₂-218, P₃-206, P₄-238 and P₅-149. A search in the mass-spectral and special pesticide libraries revealed that P₁-140, P₂-218, and P₃-206 are 4-fluorobenzoic acid, (2-fluorophenyl) (4-fluorophenyl) methanone, and 1-(4-fluorophenyl)-2- (1H-1,2,4-triazol-1-yl) ethanone, respectively. And P₄-238 is the dihydroxylated products of P₃-206. The intermediate P₄-238 has two structural isomers, and it is not possible to distinguish the two based on the acquired data. Despite a thorough search, we failed to identify the structure

of P₅-149, and it is not possible to deduce a reasonable structure based on its MS fragments.




Fig. 11. TIC chromatogram from HPLC/MS/MS




Fig. 12. Proposed transformation pathways of flutriafol in the $Ag^+/S_2O_8^{2-}$ system

In the HPLC/MS/MS analyzed (Fig. 11), the intermediates P₂-218, P₃-206 and P₄-238 are also found. Additionally, there are P₆-318 and P₇-334 that are ascribable to mono- and dihydroxylated flutriafol, respectively. The hydroxylation products were reported before by Ghauch [27]. It is noted that P₇-334 is more polar than P₆-318, in agreement

with the fact that the former is a di-hydroxylated product while the latter mono-hydroxylated product of flutriafol.

Based on the degradation intermediates and the contribution of ROS, three degradation pathways are proposed as illustrated in Fig. 12. There is the $\cdot\text{SO}_4^-$ attack toward the carbon atom (C8) connected to the tertiary alcohol, and 4-fluorobenzoic acid (P_1 -140) and (2-fluorophenyl) (4-fluorophenyl) methanone (P_2 -218) are produced as a result of C–N bond cleavage. On the other hand, the C–C bond (C₇–C₉) of tertiary alcohol is broken also due to $\cdot\text{SO}_4^-$ attack to give 1-(4-fluorophenyl)-2-(1H-1,2,4-triazol-1-yl) ethanone (P_3 -206) and its dihydroxylated products (P_4 -238). Furthermore, $\cdot\text{OH}$ reacts with the flutriafol molecule to give the monohydroxylated and dihydroxylated products.

3.5. ENVIRONMENTAL IMPLICATION

Based on the above results, the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system can be used as an in situ oxidation method to remove flutriafol in waste-water treatment. However, attention has to be paid for the optimization of reaction parameters such as pH values and Ag^+ and $\text{S}_2\text{O}_8^{2-}$ concentrations. The results of the present studies provide insights into the transformation and degradation of flutriafol under the influence of $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$. If the degradation intermediates were toxic, undesired ill effects would be resulted. It is hence essential to determine whether it is a necessity to achieve complete mineralization of degradation intermediates during the oxidation process. With the degradation pathways clarified and the intermediates identified, it is possible to examine the toxicity of the overall degradation process and have better understanding on its suitability in environment protection.

3. CONCLUSIONS

Flutriafol can be degraded efficiently in the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system by $\cdot\text{SO}_4^-$ which is generated when $\text{S}_2\text{O}_8^{2-}$ is activated by Ag^+ . It was observed that an increase in the concentration of either Ag^+ or $\text{S}_2\text{O}_8^{2-}$ in the system results in enhancement of flutriafol degradation efficiency. Also, a decrease of flutriafol concentration results in increase of degradation rate. Furthermore, the results of kinetics studies indicate that chelating agents, EDTA- Na_2 , $\text{Na}_2\text{S}_2\text{O}_3$ and dextrose monohydrate (DM), have negative effect on the performance of the $\text{Ag}^+/\text{S}_2\text{O}_8^{2-}$ system due to their ability in capturing Ag^+ . And the best degradation efficiency of flutriafol is observed at pH 3.0. Both $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$ radicals are involved in the degradation, with $\cdot\text{SO}_4^-$ contributing more. With the identification of degradation intermediates by GC/MS and HPLC/MS/MS, three degradation pathways that involve the cleavage of C–N and C–C bonds and hydroxylation of the parent compound are proposed. Comparing the oxidation ability of $\cdot\text{SO}_4^-$ and $\cdot\text{OH}$, the

cleavage of the C–N and C–C bond is a result of [•]SO₄⁻ attack, while the hydroxylation reactions are due to [•]OH. With the degradation pathways clarified, it is envisaged that the toxic effects of the degradation intermediates can be investigated and documented.

ACKNOWLEDGMENTS

The authors thank the Provincial Natural Science Foundation of Hunan (2018JJ2079, 2015JJ3056, 2015JJ2042), Scientific Research Fund of Hunan Provincial Education Department (17B061, 14K030, 13A133).

REFERENCES

- [1] WHITE P.M., POTTER T.L., CULBREATH A.K., *Fungicide dissipation and impact on metolachlor aerobic soil degradation and soil microbial dynamics*, *Sci. Total Environ.*, 2010, 408 (6), 1393.
- [2] ZHANG Q., HUA X.D., SHI H.Y., LIU J.S., TIAN M.M., WANG M.H., *Enantioselective bioactivity, acute toxicity and dissipation in vegetables of the chiral triazole fungicide flutriafol*, *J. Hazard. Mater.*, 2015, 28, 465.
- [3] YU P.Z., JIA C.H., SONG W.C., LIU F.M., *Dissipation and residues of flutriafol in wheat and soil under field conditions*, *B, Environ. Contam. Tox.*, 2012, 89 (5), 1040.
- [4] MAVROEIDI V.I., SHAW M.W., *Sensitivity distributions and cross-resistance patterns of Mycosphaerella graminicola to fluquinconazole, prochloraz and azoxystrobin over a period of 9 years*, *Crop. Prot.*, 2005, 24 (3), 259.
- [5] HERMANSON M.H., ISAKSSON E.H., TEIXEIRA C., MUIR D.C.G., COMPHER K.M., LI Y.F., IGARASHI I., KAMIYAMA K., *Current-use and legacy pesticide history in the Austfonna ice cap, Svalbard, Norway*, *Environ. Sci. Technol.*, 2005, 39 (21), 8163.
- [6] CASARA K.P., VECCHIATO A.B., LOURENCETTI C., PINTO A.A., DORES E.F.G.C., *Environmental dynamics of pesticides in the drainage area of the Sao Lourenco River Headwaters, Mato Grosso State, Brazil*, *J. Brazil. Chem. Soc.*, 2012, 23 (9), 1719.
- [7] MUNIER-LAMY C., BORDE O., *Effect of a triazole fungicide on the cellulose decomposition by the soil microflora*, *Chemosphere*, 2000, 41 (7), 1029.
- [8] MUIR D.C., TEIXEIRA C., WANIA F., *Empirical and modeling evidence of regional atmospheric transport of current-use pesticides*, *Environ. Toxicol. Chem.*, 2004, 23 (10), 2421.
- [9] GILBERT-LOPEZ B., GARCIA-REYES J.F., MOLINA-DIAZ A., *Determination of fungicide residues in baby food by liquid chromatography-ion trap tandem mass spectrometry*, *Food Chem.*, 2012, 135 (2), 780.
- [10] ZHANG Q., HUA X., YANG Y., YIN W., TIAN M., SHI H., WANG M., *Stereoselective degradation of flutriafol and tebuconazole in grape*, *Environ. Sci. Pollut. Res. Int.*, 2015, 22 (6), 4350.
- [11] YANG H., WEI H.Q., HU L.T., LIU H.J., YANG L.P., AU C.T., YI B., *Mechanism for the photocatalytic transformation of s-triazine herbicides by center dot OH radicals over TiO₂*, *Chem. Eng. J.*, 2016, 300, 209.
- [12] YANG H., AN T., LI G., SONG W., COOPER W.J., LUO H., GUO X., *Photocatalytic degradation kinetics and mechanism of environmental pharmaceuticals in aqueous suspension of TiO₂. A case of β-blockers*, *J. Hazard. Mater.*, 2010, 179, 834.
- [13] ZHANG R., SUN P., BOYER T.H., ZHAO L., HUANG C.H., *Degradation of pharmaceuticals and metabolite in synthetic human urine by UV, UV/H₂O₂, and UV/PDS*, *Environ. Sci. Technol.*, 2015, 49 (5), 3056.
- [14] LUTZE H.V., KERLIN N., SCHMIDT T.C., *Sulfate radical-based water treatment in presence of chloride: formation of chlorate, inter-conversion of sulfate radicals into hydroxyl radicals and influence of bicarbonate*, *Water Res.*, 2015, 72, 349.

- [15] YANG Y., PIGNATELLO J.J., MA J., MITCH W.A., *Comparison of halide impacts on the efficiency of contaminant degradation by sulfate and hydroxyl radical-based advanced oxidation processes (AOPs)*, Environ. Sci. Technol., 2014, 48 (4), 2344.
- [16] OH W.D., LUA S.K., DONG Z., LIM T.T., *Performance of magnetic activated carbon composite as peroxymonosulfate activator and regenerable adsorbent via sulfate radical-mediated oxidation processes*, J. Hazard. Mater., 2015, 2, 841.
- [17] ANIPSITAKIS G.P., DIONYSIOU D.D., *Radical generation by the interaction of transition metals with common oxidants*, Environ. Sci. Technol., 2004, 38 (13), 3705.
- [18] ROMERO A., SANTOS A., VICENTE F., GONZALEZ C., *Diuron abatement using activated persulphate. Effect of pH, Fe(II) and oxidant dosage*, Chem. Eng. J., 2010, 162 (1), 257.
- [19] ZHOU L., ZHENG W., JIA Y., ZHANG J., ZENG C., ZHANG Y., WANG Q., YANG X., *Ferrous-activated persulfate oxidation of arsenic(III) and diuron in aquatic system*, J. Hazard. Mater., 2013, 263, 422.
- [20] TAN C.Q., GAO N.Y., DENG Y., AN N., DENG J., *Heat-activated persulfate oxidation of diuron in water*, Chem. Eng. J., 2012, 203, 294.
- [21] MENDEZ-DIAZ J., SANCHEZ-POLO M., RIVERA-UTRILLA J., CANONICA S., VON GUNTEN U., *Advanced oxidation of the surfactant SDBS by means of hydroxyl and sulphate radicals*, Chem. Eng. J., 2010, 163 (3), 300.
- [22] ZOU J., MA J., CHEN L., LI X., GUAN Y., XIE P., PAN C., *Rapid acceleration of ferrous iron/peroxymonosulfate oxidation of organic pollutants by promoting Fe(III)/Fe(II) cycle with hydroxylamine*, Environ. Sci. Technol., 2013, 47 (20), 11685.
- [23] JI Y., FERRONATO C., SALVADOR A., YANG X., CHOVELON J.M., *Degradation of ciprofloxacin and sulfamethoxazole by ferrous-activated persulfate. Implications for remediation of groundwater contaminated by antibiotics*, Sci. Total Environ., 2014, 47, 2800.
- [24] ZENG J., YANG H., LIU H., YI X., YANG L., YI B., *Common characteristic assessments of transformation mechanism for substituted phenylurea herbicides by reactive oxygen species (ROSs) during photocatalytic process*, Chem. Eng. J., 2015, 273, 519.
- [25] YANG H., LIU H., HU Z., LIANG J., PANG H., YI B., *Consideration on degradation kinetics and mechanism of thiamethoxam by reactive oxidative species (ROSs) during photocatalytic process*, Chem. Eng. J., 2014, 245 (1), 24.
- [26] YANG H., YANG L., HU Z., LIU H., PANG H., YI B., *General transformation law of chlorinated acetanilide herbicides by reactive oxidative species (ROSs) during photocatalytic process*, J. Water Process. Eng., 2015, 8, e66.
- [27] GHAUCH A., *Rapid removal of flutriafol in water by zero-valent iron powder*, Chemosphere, 2008, 7, 1816.